



	Turf: Seed	Turf: New Sod	Turf: Established	Plantings: Ball & Burlap or Container	Plantings: Soil Injection	Transplants	Compost Tea
MycoApply® Ultrafine Endo (Suspendable Powder)	Mix with seed at a rate of 0.5 ounce per 1,000 square feet or 15 grams per 100 square meters. Can be blended with fertilizer.	Apply during installation of turf at a rate of 0.67 to 1.62 ounces per 1,000 square feet or 20-50 grams per 100 square meters. Can be blended with fertilizer.	Apply as a broadcast during aerification of turf at a rate of 0.67 to 1.62 ounces per 1,000 square feet or 20-50 grams per 100 square meters. If multiple applications are made in a year, then reduce the rate by 40% in subsequent applications. Irrigate well to ensure movement into root zone for established turf. The product can be blended with fertilizer, if desired.	Incorporate 0.5 oz with 2 cubic feet of backfill soil (2.5 grams per 10 liters) or make a water suspension by mixing 2 oz in 5 gallons of water (3 grams per liter) and apply sufficient suspension to moisten the sides of the rootball and the backfill soil.	Prepare a suspension using 1 pound per 20 gallons (3 grams per liter) and inject into multiple holes near the dripline of the trees or shrubs.	Dip plugs in a suspension of 2 oz per 5 gallons of water (3 g /liter) for 10-15 seconds to obtain good contact with substrate. For bare-rooted transplants, a spreader sticker may be added to the dip rate suspension, and the roots should be thoroughly sprayed with the suspension.	Use 1 pound per 50 gallons of tea (2.4 grams per liter) as a soil drench. Apply tea at a normal rate. Water in after application to ensure propagule movement into the root zone.
	Turf: Seed	Turf: New Sod	Turf: Established	Plantings: Ball & Burlap or Container	Plantings: Furrow or Broadcast	Transplants	
MycoApply® Endo (Granular)	Mix with seed at a rate of 1 ounce per 1,000 square feet or 30 grams per 100 square meters. Can be blended with fertilizer.	Apply at a rate of 0.1 -0.25 pounds per 1,000 square feet or 50-125 grams per 100 square meters. Can be blended with fertilizer.	Apply as a broadcast during aerification of turf at a rate of 0.1 - 0.25 pounds per 1,000 square feet or 50-125 grams per 100 square meters. If multiple applications are made in a year, then reduce the rate by 40% in subsequent applications. Irrigate well to ensure movement into root zone for established turf. The product can be blended with fertilizer, if desired.	Incorporate 1 oz with 2 cubic feet of backfill soil (5 grams per 10 liters) or make a water suspension by mixing 4 oz in 5 gallons of water (6 grams per liter) and apply sufficient suspension to saturate the sides of the rootball and the backfill soil.	Furrows: Best to mix with seed or fertilizer at the rate of 0.5 oz per 1,000 linear feet (1 gram per 20 linear meters). Broadcast: Apply 0.25 - 0.35 pounds per 1,000 square feet or 1.25-1.7 grams per square meter.	Dip plugs in a suspension of 4 oz per 5 gallons of water (6 grams/liter) for approximately 10-15 seconds to obtain good contact with roots and substrate. For bare-rooted transplants, a spreader sticker may be added to the dip rate suspension, and the roots should be thoroughly sprayed with the suspension.	

MycoApply Products are compatible with a variety of fungicides. Check our website or contact us for a full compatibility list before applying a fungicide.

	Plantings: Ball & Burlap or Container	Plantings: Soil Injection	Transplants	Compost Tea
MycoApply® Ultrafine Endo/Ecto (Suspendable Powder)	Incorporate 0.5 oz with 2 cubic feet of backfill soil (3 grams per 10 liters) or make a water suspension by mixing 2 oz in 5 gallons of water (3 grams per liter) and apply sufficient suspension to moisten the sides of the rootball and the backfill soil.	Prepare a suspension using 1 pound per 20 gallons (6 grams per liter) and inject into multiple holes near the dripline of the trees or shrubs.	Dip plugs in a suspension of 2 oz per 5 gallons of water (3 g /liter) for 10-15 seconds to obtain good contact with substrate. For bare-rooted transplants, a spreader sticker may be added to the dip rate suspension, and the roots should be thoroughly sprayed with the suspension.	Use 1 pound per 50 gallons of tea (2.4 grams per liter) as a soil drench. Apply tea at a normal rate. Water in after application to ensure propagule movement into the root zone.
MycoApply® Soluble MAXX (Suspendable Powder)	Incorporate 0.5 oz with 2 cubic feet of backfill soil (3 grams per 10 liters) or make a water suspension by mixing 2 oz in 5 gallons of water (3 grams per liter) and apply sufficient suspension to saturate the sides of the rootball and the backfill soil.	Prepare a suspension using 1 pound per 50 gallons (2.4 grams per liter) and inject into multiple holes near the dripline of the trees or shrubs.	Dip plugs in a suspension of 2 oz per 5 gallons of water (3 g /liter) for 10-15 seconds to obtain good contact with substrate. For bare-rooted transplants, a spreader sticker may be added to the dip rate suspension, and the roots should be thoroughly sprayed with the suspension.	Use 1 pound per 50 gallons of tea (2.4 grams per liter) as a soil drench. Apply tea at a normal rate. Water in after application to ensure propagule movement into the root zone.

	Plantings: Ball & Burlap or Container	Plantings: Furrow or Broadcast	Transplants
MycoApply® Endo/Ecto (Granular)	Incorporate 1 oz with 2 cubic feet of backfill soil (5 grams per 10 liters) or make a water suspension by mixing 4 oz in 5 gallons of water (6 grams per liter) and apply sufficient suspension to saturate the sides of the rootball and the backfill soil.	Furrows: Best to mix with seed or fertilizer at the rate of 0.5 oz per 1,000 linear feet (1 gram per 20 linear meters). Broadcast: Apply 0.25 - 0.35 pounds per 1,000 square feet or 1.25-1.7 grams per square meter.	Dip plugs in a suspension of 4 oz per 5 gallons of water (6 grams /liter) for approximately 10-15 seconds to obtain good contact with roots and substrate. Keep suspension agitated. For bare-rooted transplants, a spreader sticker may be added to the dip rate suspension, and the roots should be thoroughly sprayed with the suspension.

	Turf: Seed	Turf: New Sod	Turf: Established
MycoApply® Injector Endo (Concentrated Suspendable Powder)	Mix concentrated powder with seed at a rate of 0.25 gram per 1,000 square feet or 0.25 gram per 100 square meters. Can be blended with fertilizer, if desired.	Apply as a liquid suspension drench during installation of turf at a rate of 0.3 to 0.8 gram per 1,000 square feet or 0.3 to 0.8 gram per 100 square meters. Can be applied via 1:100 injector at a rate of 20 grams per 100 gallons, using drip, spray, or sprinkler irrigation. Can be blended with fertilizer, if desired.	Apply as a liquid suspension drench during aerification of turf at a rate of 0.3 to 0.8 gram per 1,000 square feet or 0.3 to 0.8 gram per 100 square meters. Can be applied via 1:100 injector at a rate of 20 grams per 100 gallons, using drip, spray, or sprinkler irrigation. The product can be blended with fertilizer, if desired.

MycoApply Products are compatible with a variety of fungicides. Check our website or contact us for a full compatibility list before applying a fungicide.

Version 1.8 – Updated 7/1/2019